Proforma of ultrasound and MRI signs in PAS

Suspected Placenta Accreta Spectrum: imaging reporting template						
Date of examination	//					
Gestational Age	weeksda	ays				
Demographics and Risk Factors						
Parity						
Number of previous CS						
Number of previous surgical evacuations (including TOP)						
	Yes	No	Not known			
Previous uterine surgery (e.g. myomectomy, endometrial ablation)?						
History of PAS?						
Was CS scar pregnancy suspected first trimester?						
Placenta praevia on US?						

Ultrasound Signs 'x' as appropriate	Yes	No	Unsure	MRI Signs 'x' as appropriate	Yes	No	Unsure
Loss of clear zone				Loss of retroplacental low T2 line			
Myometrial thinning				Myometrial thinning (<1mm)			
Uterine bulge				Uterine bulge			
Focal exophytic placental mass – through serosa, parametrium				Focal exophytic placental mass – through serosa, parametrium			
Bladder wall interruption				Bladder wall interruption			
Lacunae				Low T2 placental bands			

Colour Doppler		Heterogeneity		
Placenta lacunae feeder vessels		Intraplacental fetal vessels (>3mm)		
Abnormal vasculature at plac- myometrial interface, serosa, bladder		Abnormal vasculature at plac-myometrial interface, serosa, bladder		

Clinical Significance of Imaging Findings	High	Intermediate	Low	
Extent of PAS	Focal		Diffuse	