

## NATIONAL GI ENDOSCOPY QI PROGRAMME KEY QUALITY INDICATORS

Key Quality Indicator	Key Quality Target	Additional Information
<b>COLONOSCOPY</b>		
Caecal intubation rate (CIR)	Minimum: $\geq 90\%$ Achievable: $\geq 95\%$	CIR is calculated based on all colonoscopies performed as Endoscopist 1 or Endoscopist 2.
Comfort Score	$\geq 90\%$	90% of colonoscopies should have a comfort score of between 1 and 3 on the Gloucester Scale.
Polyp detection	$\geq 20\%$	N/A
Bowel Preparation	Minimum: $\geq 90\%$ Achievable: $\geq 95\%$	N/A
<b>OESOPHAGOGASTRODUODENOSCOPIES (UPPER GI)</b>		
Duodenal 2nd part intubation (Duo 2)	$\geq 95\%$	N/A
Retroflexion	$\geq 95\%$	N/A
<b>SEDATION</b>		
Midazolam	<p><b>Patients Aged below 70 years:</b> Median dose is <math>\leq 5</math> mg per endoscopist</p> <p><b>Patients Aged above 70 years</b> Median dose is <math>\leq 3</math> mg per endoscopist</p>	This KQI applies to both colonoscopies and OGDs.